

Relationships between extreme flows and microbial contamination in inland recreational swimming areas

Kathryn Vincent^a, Danswell Starrs^a, Victoria Wansink^a, Natasha Waters^a and Aparna Lal ^{b,*}

^a Australia

^b Australian National University, Canberra, Australia

*Corresponding author. E-mail: aparna.lal@anu.edu.au

 AL, 0000-0001-7062-1468

ABSTRACT


Inland recreational swimming sites provide significant social value globally. This study focused on public recreational swimming sites across the Murrumbidgee River and its tributaries in the Australian Capital Territory (ACT) throughout the swimming season (September–April) from 2009 to 2020 to determine whether high intestinal enterococci concentrations could be predicted with flow exceedance and routinely monitored physical and chemical parameters of water quality. Enterococci concentrations were positively correlated with the turbidity associated with high-flow conditions. The predictive accuracy of high enterococci levels during high-flow conditions was good (mean percentage correctly classified, 60%). The prediction of high enterococci levels at low flows was significantly less reliable (mean percentage correctly classified, 12–15%). As the ACT is expected to experience decreases in rainfall overall but increases in extreme rainfall events due to climate change, understanding the drivers of elevated intestinal enterococci under extreme flow conditions remains important from a public health perspective.

Key words: Australia, extreme flows, inland rivers, primary contact, public health, swimming

HIGHLIGHTS

- Climate change is anticipated to increase the frequency and magnitude of extreme rainfall.
- Extreme riverine flows were associated with high enterococci concentration.
- High flows and turbidity were good predictors of high enterococci concentrations.
- High enterococci during low flows were not well explained by the water quality or flow.

GRAPHICAL ABSTRACT



1. INTRODUCTION

The quality of water in naturally occurring freshwater systems is under increasing pressure from development and population expansion (Buckerfield *et al.* 2019). These impacts will likely be exacerbated by extreme weather events associated with climate change (Alkolibi 2002; Kistemann *et al.* 2002; Signor *et al.* 2005; Sidhu *et al.* 2012). Extreme precipitation can substantially alter the risks to public health, either due to mobilisation of contaminants during flooding or of higher concentrations due to reduced flow volumes during prolonged dry periods (Joosten *et al.* 2017; Buckerfield *et al.* 2019). Heavy rainfall driving the flushing of contaminants into surface and groundwater sources is a well-established precursor to increased gastrointestinal illness (Curriero *et al.* 2001; Tornevi *et al.* 2013). Positive associations between enterococci concentrations and gastrointestinal illness in swimmers (Boehm & Sassoubre 2014) and high-flow conditions have frequently been reported in the literature (Chu *et al.* 2014; Strauch 2017). While there is general consensus on the impact of high-flow events in freshwater systems on public health, the effect of low-flow conditions following prolonged dry periods on the health risks is not well assessed.

Enterococci that are found in human and animal faeces are frequently used as faecal indicator bacteria (FIB) as a proxy for water quality and to assess the public health risks in recreational waters (Boehm & Sassoubre 2014). Four years of data in a Mediterranean coastal river in low-flow conditions showed that bacteria were stored in fine riverbed sediments, creating an in-stream store of bacteria (Chu *et al.* 2014). The presence of submerged aquatic vegetation provides conditions conducive to higher mean densities of enterococci than in sediment or water (Badgley *et al.* 2010), indicating another potential source of enterococci contamination and/or persistence during low-flow conditions. In Florida, USA, beach management activities such as reducing human, dog and bird densities, and reducing homeless populations were associated with lower enterococci concentrations (Kelly *et al.* 2018) suggesting that non-point sources of pollution may be an important source of contamination during low-flow conditions. Collectively, results from previous work identify a clear need to better understand how extreme high- and low-flow conditions can alter the water-related risks to public health.

In Australia, the majority of reported outbreaks of water-related gastrointestinal illness are related to recreational use (Dale *et al.* 2010; Sidhu *et al.* 2012), with concentrations of FIB in urban stormwater runoff frequently exceeding acceptable limits after storm events. Future climate projections for rainfall in Australia include a decrease in overall rainfall in southern Australia, but an increase in intense rainfall events throughout the country (IPCC 2014; BOM & CSIRO 2018). Being able to

predict the levels of microbial pollution has locally relevant implications for current management actions, as well as planning for increased risk associated with water-based recreation due to climate change.

The current study focuses on the Australian Capital Territory (ACT). In the ACT, less rain is projected during the winter and spring seasons by 2090, and daily rainfall intensity is expected to increase (Webb & Hennessy 2015). The main aim of this study is to examine and predict how flow and flow variability is associated with intestinal enterococci concentration, while considering additional water quality parameters that are important for management of natural swimming areas. Specific objectives were to (i) quantify flow exceedance at multiple swimming sites over a long time period, (ii) quantify relationships between intestinal enterococci concentrations, flow exceedance, and water quality parameters including turbidity, electrical conductivity, and water temperature, and (iii) predict intestinal enterococci concentrations during extreme flow (high- and low-flow conditions). Additionally, we establish a link between visitation to these areas using traffic count data and periods of high intestinal enterococci concentration.

2. METHODS

2.1. Study area

Within the ACT, the Murrumbidgee River and its riparian corridor are managed primarily for conservation purposes, and as a place of recreation for residents of the ACT and surrounding regions. Currently, there are seven designated recreational swimming sites along the Murrumbidgee River, which flows through the ACT for approximately 66 km. Two additional designated primary recreational swimming sites occur on the Paddys River (which is a tributary of the Cotter River) and the Cotter River, which is a tributary of the Murrumbidgee River (Figure 1). The Murrumbidgee River catchment upstream of the ACT is predominantly rural, with livestock grazing and cropping being the predominant land-uses adjacent to the Murrumbidgee River (ABARES 2021). A considerable proportion of the Paddys River catchment includes agricultural land-uses as well.

2.2. Data sources and calculations and flow exceedance analysis

2.2.1. Intestinal enterococci concentration data

Intestinal enterococci data were obtained from the ACT Data's 'Open Data Portal' (ACT Government 2019), which is collected and analysed by the ACT Health Protection Service weekly (Monday or Tuesday, with results reported on Friday) during the open swimming season (September–April). Data were from Austral Spring 2009 to Austral Autumn 2020. The intestinal enterococci counts are obtained through analysis by the ACT Government Analytical Laboratory (ACTGAL), a NATA-accredited facility. These counts are reported as the number of intestinal enterococci organisms or colony-forming units (CFU) per 100 mL of sampled water. According to the ACT Guidelines for Recreational Water Quality 2014, if intestinal enterococci counts are >200 CFU/100 mL another sample is taken. If the repeat sample is >200 CFU/100 mL then the site is advised as closed for primary contact recreation, and the relevant territory bodies are advised to facilitate communication of the risk to the public, with signage displayed (ACT Government Health 2014). To reopen a site, two consecutive samples must produce a count ≤200 CFU/mL. Prior to 2014, a single sample >200 CFU/100 mL was required to advise the closing of a site for primary contact recreation. As such, results are reported as the advisory status for sites, rather than closed, as other water quality and management issues (e.g. blue-green algae) may lead to a site being closed in addition to intestinal enterococci count data.

2.2.2. Flow exceedance calculation and water quality data

River streamflow gauges for the Murrumbidgee River, Gudgenby River, Tuggeranong Creek, Paddys River, and Cotter River were used to explore the role of streamflow on intestinal enterococci concentrations. Daily streamflow data (mL/day) from 1980 to 2020 were used to calculate the long-term streamflow exceedance curves. Summation of streamflow from gauges on tributaries with the Murrumbidgee River gauges was used to compute streamflows for intestinal enterococci sampling sites not associated with a streamflow gauge. This produced five unique streamflow datasets for analysis (Supplementary Material, Figure 1). Missing streamflow data were interpolated from upstream or downstream gauges, or from time series of streamflow data generated by a calibrated catchment model for the ACT (eWater Source model). Flow exceedance was calculated as:

$$P = 100 \times \left[\frac{M}{n + 1} \right] \quad (1)$$




Figure 1 | Location of study sites along the Murrumbidgee River and its tributaries in the ACT, Australia.

where P is the probability that a given discharge value will be equalled or exceeded; M is the ranking of the discharge value; n is the number of values in the dataset.

Analyses were based on the flow exceedance probability (FEP) rather than daily streamflow, to account for the differences in streamflow between sites which make between-site comparisons problematic. Analyses focussed on exploring the relationship between intestinal enterococci count (CFU/100 mL) and FEP at each intestinal enterococci sampling location. Intestinal enterococci counts at Cotter Campground were examined with both the Cotter River flow and Paddys River flow to assist in determining whether tributary inputs from the Paddy's River better explain intestinal enterococci counts at the Cotter River swimming site. To further understand the possible role of antecedent flow conditions (e.g. 1–7 days prior to intestinal enterococci samples being taken), flow exceedance was lagged sequentially each day for up to 7 days. Antecedent flow variability was calculated as the difference between FEP and lagged FEP (daily, up to 7 days), divided by FEP. Antecedent flow variability was $\log_{10}(x + 1)$ transformed prior to analysis to minimise

the leverage due to outlier values.

$$V = \frac{FEP_i - FEP_0}{FEP_0}$$

where i is between 1 and 7 days preceding, and 0 is the day on which the intestinal enterococci sample was collected. V is the antecedent flow variability; FEP is the flow exceedance probability.

A single *in situ* water quality sonde is located on the Murrumbidgee River at the upstream margin of the ACT (Lobbs Hole, gauge #410761; Figure 1). This sonde records water quality parameters (turbidity, water temperature, dissolved oxygen, pH, and electrical conductivity) at 15-min intervals. Time-weighted mean daily values were calculated for each parameter for analysis. Missing data (e.g. due to equipment malfunction) were not infilled or replaced, as no alternate data source was available. Water quality data were standardised (centred and scaled to mean = 0 and standard deviation = 1) both within year, and within month and year to remove annual and seasonal patterns from the data. Given the extreme variability in mean daily turbidity, turbidity data were also categorised into 'low' and 'high' turbidity, based on the recommended threshold value (25 NTU) presented in the ANZECC guidelines for upland rivers in south-eastern Australia (ANZECC & ARMICANZ 2000).

2.2.3. Traffic count data

Road traffic count data were obtained from ACT Parks and Conservation Service for the Murrumbidgee River corridor. Vehicular traffic into carparks at some of the swimming sites has been monitored periodically since 2011. Data are collected from traffic counters on the first day of each month. Where there were missing data (due to equipment malfunction, theft, or vandalism) no attempt was made to infill or replace the missing data, as no reference data were available.

2.3. Statistical analysis

Relationships between \log_{10} transformed intestinal enterococci concentrations (CFU/100 mL), water quality parameters (both raw and standardised), and streamflow characteristics, including flow exceedance and antecedent flow exceedance variability were examined. Intestinal enterococci concentrations were $\log_{10}(x + 1)$ transformed due to non-normally distributed modelled residuals, which were adequate after transforming the response variable. Intestinal enterococci count data for the three sites closest to the Lobbs Hole streamflow gauge (Tharwa Bridge, Point Hut, and Pine Island) were examined for relationships with water quality parameters and the above-mentioned metrics for streamflow. Intestinal enterococci concentration and streamflow metrics were examined for every intestinal enterococci monitoring site. No attempt was made to apply the water quality data to other intestinal enterococci monitoring sites, as additional significant tributary inputs (including urban inflows) may lead to incorrect interpretations. A single model with all explanatory variables was fitted for each site to examine sources of variation in intestinal enterococci concentrations.

Tests for significant relationships ($\alpha = 0.05$) between \log_{10} transformed intestinal enterococci concentration and streamflow and water quality were performed by linear regression. Models were fitted with generalised least squares (GLS) and an AR(1) autoregressive error structure to account for temporal autocorrelation in residuals. Explanatory variables were examined for collinearity prior to fitting. Suitability of model fit was examined for all fitted models, through examination of model residuals and AIC values.

To inform whether the prediction of high intestinal enterococci could be undertaken using streamflow and/or water quality data to inform future management of the public health risks, a repeated k -folds cross-validation analysis was conducted to test the efficacy of a linear model to predict high (>200 CFU/100 mL) intestinal enterococci concentrations at each monitoring location. Intestinal enterococci data were converted to a binary variable, around the management threshold of 200 CFU/100 mL. The dataset was split into five randomly selected blocks, with four used for training and one for validation. Each block was sequentially used as the validation set, with the remaining used for training the model. This was repeated 100 times (with data randomly assigned to blocks after each iteration) to evaluate model performance. Variability in model performance was explored, and residual risk to the community was inferred through relative changes in recreational water quality. Monthly traffic counter (2011–2020) data for sites where data were available were standardised by site and plotted against the proportion of intestinal enterococci samples exceeding 200 CFU/100 mL per month to visually examine any potential correlation between visitation rates and increased risk of exposure to high intestinal enterococci concentrations.

3. RESULTS

3.1. Intestinal enterococci concentrations varied both spatially and temporally

Intestinal enterococci count data (CFU/100 mL) varied spatially across the nine river swimming sites sampled (Figure 2). Temporal variation was seen with both annual and seasonal patterns (Figures 2 and 3). Murrays corner (Paddys River) exhibited the highest (mean \pm standard error (SE)) intestinal enterococci counts over the period of study (436 ± 80), with Kambah Pool exhibiting the lowest mean intestinal enterococci counts (118 ± 15). February had the highest mean intestinal enterococci counts (426 ± 83), while September had the lowest counts (42 ± 9). February had the highest percentage (nearly 34%) of occasions where the CFU was >200 CFU/100 mL (Supplementary Material, Table 1).

Advised periods of no primary contact recreation varied through time and between swimming sites (Figure 3). This was further compounded by the change in ACT Guidelines for Recreational Waters, that occurred between the 2013/14 and 2014/2015 swimming seasons. Casuarina Sands exhibited the lowest average number of days advised closed to primary contact recreation (11.36 ± 3.67), while Murrays Corner on the Paddy's River exhibited the highest average number of days advised to be closed (104.91 ± 24.12) (Figure 3).

3.2. Intestinal enterococci varied with flow duration, antecedent flow exceedance variability, and turbidity

At high levels of turbidity, there was an inverse association between FEP and enterococci counts (Figure 4). This relationship held at all sites, except Paddys River, Cotter River, and Casuarina Sands which exhibited no significant relationship between flow exceedance and intestinal enterococci counts (Supplementary Material, Table 2). Inclusion of Paddys River flow to explain variation in intestinal enterococci counts in the Cotter River provided a better model fit than using the Cotter River discharge (AIC: 415.69 vs. 433.52). An examination of public visitation to swimming sites suggests there is a strong seasonal pattern in traffic counts with the largest standardised counts occurring in December, and the lowest in June (Figure 5).

3.3. Model performed well at predicting high-flow intestinal enterococci, but poorly at low flow

The inclusion of water quality parameters further revealed that high turbidity associated with high-flow periods is positively associated with increased intestinal enterococci counts, whereas high flow associated with lower turbidity was not associated




Figure 2 | Box-and-whisker plots of intestinal enterococci count (CFU/100 mL) by month for nine river swimming locations in the ACT 2009–2019, on log₁₀ scale. The red dashed line indicates the management threshold of 200 CFU/100 mL, over which swimming locations are advised as closed to primary contact recreation due to intestinal enterococci counts. See Figure 1 for map of sampling locations. Please refer to the online version of this paper to see this figure in colour: <http://dx.doi.org/10.2166/wh.2022.294>.




Figure 3 | Putative number of days each site was advised as closed to primary contact recreation due to intestinal enterococci counts per year from Spring 2009 to Autumn 2020. Note that year is the water year (July–June), with the swimming period defined as September–April.

Table 1 | Model summary showing the prediction classification accuracy (proportion of samples correctly classified as >200 CFU/100 mL, mean and range) for all data, proportion correctly classified as >200 CFU/100 mL under high flow (FEP < 50), proportion correctly classified as >200 CFU/100 mL under low flow (FEP > 50), and RMSE for fitted models

Site	All data	High flow	Low flow	RMSE (all data)
Tharwa Bridge	0.82 (0.80–0.83)	0.60 (0.54–0.68)	0.12 (0.06–0.16)	0.52 (0.51–0.54)
Point Hut	0.85 (0.83–0.86)	0.59 (0.53–0.67)	0.13 (0.05–0.2)	0.49 (0.48–0.51)
Pine Island	0.85 (0.83–0.86)	0.59 (0.53–0.64)	0.14 (0.05–0.25)	0.49 (0.48–0.51)
Kambah Pool	0.86 (0.84–0.87)	0.33 (0.29–0.40)	0.07 (0–0.15)	0.47 (0.46–0.49)
Casuarina Sands	0.88 (0.87–0.88)	0.39 (0.36–0.44)	0.16 (0.08–0.17)	0.46 (0.45–0.49)
Uriarra Crossing	0.87 (0.86–0.87)	0.40 (0.35–0.46)	0.18 (0.10–0.20)	0.45 (0.44–0.47)
Swamp Creek	0.77 (0.76–0.79)	0.49 (0.41–0.56)	0.16 (0.13–0.21)	0.52 (0.52–0.53)
Paddys River	0.70 (0.68–0.71)	0.71 (0.67–0.75)	0.60 (0.55–0.62)	0.42 (0.41–0.45)
Cotter River	0.79 (0.78–0.80)	0.29 (0.23–0.32)	0.04 (0.02–0.07)	0.42 (0.41–0.43)
Cotter River/Paddys Flow	0.81 (0.80–0.82)	0.49 (0.45–0.52)	0.004 (0–0.04)	0.40 (0.39–0.42)

Mean and range calculated from 100 iterations of a k -folds cross-validation ($k = 5$).

with higher intestinal enterococci counts. Antecedent flow conditions up to 7 days prior were associated with intestinal enterococci counts, with increased variability in antecedent flow exceedance being associated with high intestinal enterococci counts (Supplementary Material, Table 2).

Linear models exploring the relationship between intestinal enterococci counts, flow exceedance, antecedent flow variability, month, turbidity, and site explained 47.9, 40.3, and 40.8% of the variation in intestinal enterococci counts at Tharwa Bridge, Point Hut, and Pine Island, respectively (Supplementary Material, Table 3). Flow exceedance, flow exceedance variability, and month were all significant predictors at all sites.

A k -folds validation produced a root mean square error (RMSE) of 0.525 (range 0.515–0.54) for Tharwa Bridge. Overall, minimal variability in RMSE indicates that models are robust (Table 1). Model prediction efficiency for high flow (flow exceedance percentile < 50) and high intestinal enterococci counts (>200 CFU/100 mL) was variable, with a range of 29.67%




Figure 4 | Relationship between \log_{10} intestinal enterococci concentration (CFU/100 mL), streamflow exceedance probability and turbidity at (a) Tharwa Bridge, (b) Point Hut Crossing, and (c) Pine Island on the Murrumbidgee River. Red dashed line denotes the management threshold of 200 CFU/100 mL. Increased intestinal enterococci concentrations at higher stream flows is associated with high turbidity (≥ 25 NTU). Please refer to the online version of this paper to see this figure in colour: <http://dx.doi.org/10.2166/wh.2022.294>.




Figure 5 | Mean number of day sites were advised as closed to primary contact recreation due to intestinal enterococci \pm SE (black) (2009–2020) by month, and standardised monthly traffic counts \pm SE (blue) for recreational swimming sites (2010–2019). Please refer to the online version of this paper to see this figure in colour: <http://dx.doi.org/10.2166/wh.2022.294>.

correct (the Cotter River with the Cotter River flow) to 71.24% correct (Paddys River). Conversely, the prediction accuracy of high intestinal enterococci under low-flow conditions (flow exceedance percentile $>50\%$) was poor, ranging from 0.47% correct (the Cotter River with the Paddys River flow) to 16.74% correct (Casuarina Sands).

4. DISCUSSION

Overall, the current study found that extreme flows (high and low) were consistently associated with high intestinal enterococci concentration. We found a consistent positive relationship between high (>200 CFU/100 mL) intestinal enterococci concentrations, turbidity equal to or greater than 25 NTU, and high-flow conditions. High flows are commonly associated with the elevated FIB concentrations. Storm runoff, land-use such as agriculture, and stormwater drain inputs have been implicated as a key driver of short-term increases of intestinal enterococci in freshwater systems (Coffey *et al.* 2013; Daly *et al.* 2013; Chu *et al.* 2014). Disturbance of riverbeds – whether through increased streamflow, direct disturbance from users or livestock, or other means – has been found to resuspend sediment and FIB within sediments (Cho *et al.* 2010b; Cho *et al.* 2016) leading to a secondary non-point source of FIB contamination. The consistent association with high turbidity during elevated flow conditions may be partly attributed to the resuspension of sediments within the stream channel during high-flow conditions, which is known to increase the pathogen and FIB levels in the water (Anderson *et al.* 1997; Gao *et al.* 2013; Abia *et al.* 2017). Suter *et al.* (2011) also found enterococci concentrations positively correlated with turbidity. Increased turbidity may also limit the amount of solar radiation penetrating the water column, decreasing its potential to inactivate microbial contaminants (Williamson *et al.* 2017). Previous studies have modelled the effect of solar irradiance on FIB under high- and low-flow conditions (Deely *et al.* 1997; Coffey *et al.* 2013; Herrig *et al.* 2019). Herrig *et al.* (2019) showed that global solar irradiance was inversely associated with the FIB concentrations. Discharge from urban land-use can cause browning of surface waters, with the potential for highly turbid conditions. Our study highlights the implications of such conditions for the public health risks associated with recreational use in such waterways (Meyer *et al.* 2005).

There is limited literature that examines the association between low-flow conditions and high intestinal enterococci levels. Increased microbial loads due to point sources during low-flow periods can occur due to reduced dilution as a result of reduced water volumes (Thilakarathne *et al.* 2018). Using 4 years of data on Mediterranean coastal river in low-flow conditions, Chu *et al.* (2014) found that bacteria were stored in riverbed sediments forming in-stream stores that were the source of elevated bacterial levels during the first flush event. In addition to bacteria persistence during low-flow conditions, the source of these bacteria may also be different. Due to point and non-point sources of FIB present in sediments, the persistence of FIB in sediments may be highly variable. Previous studies in Australia's Murray Darling Basin have shown that when relative discharge is controlled for, water temperature is positively associated with high cyanobacteria abundance (Lal & Hargreaves 2020). Therefore, high water temperature associated with low flows may also partly explain the high enterococci levels.

The inclusion of water quality parameters (namely, turbidity) improved the ability to model and predict high intestinal enterococci concentrations. Numerous studies previously have demonstrated that both external loading from catchments, in addition to internal resuspension of FIB within waterways during high-flow events can lead to increased FIB concentrations (Jamieson *et al.* 2005; Cho *et al.* 2010a; Walters *et al.* 2014; Ribolzi *et al.* 2016). Across aquatic systems globally, increased turbidity is commonly associated with intensification of land-use and poor riparian zone condition and management (Smith *et al.* 2013; Terry *et al.* 2014; Davis *et al.* 2015). Within the upper Murrumbidgee River catchment, gully erosion has been historically problematic, however, this is now considered to be less so than previously (Olley & Wasson 2003). Conversely, poor riparian condition evidenced by the loss of riparian zone depth and complexity (O'Reilly *et al.* 2021) and access by livestock to watercourses may be a significant source of both FIB and continued deposition of sediments to waterways. There is potential for riparian zone restoration, fencing of livestock out of riparian zones to contribute to improved riparian zone condition, and reduced inputs of FIB and sediments to the Murrumbidgee River, leading to overall improved waterway condition and reduced risk to public health.

4.1. Study limitations, management implications, and further work

Results from this study highlight the ability to predict intestinal enterococci concentrations using routinely collected hydrological and water quality parameters. A strength of this study lies in the consistently high accuracy of prediction with high-flow conditions, suggesting that water quality parameters could be used alongside flow data for other sites to predict intestinal enterococci concentration. Equally important is the consistent finding of low prediction accuracy of high intestinal enterococci concentration at low-flow conditions, indicating the potential importance of non-point sources of contamination, including humans. Collectively, study findings provide a clearer distinction of the potential drivers and therefore management options for reducing intestinal enterococci concentration at high- and low-flow conditions across multiple sites using routinely collected data.

Timeliness is a key metric when assessing the effectiveness of a public health response. Our results highlight the potential impact of the time delay between sample collection and the reporting of results. In the current study, results of samples collected on a Monday/Tuesday were only reported later in the week. Due to this, our analyses were conducted at the weekly scale. The strong association between high flow and high enterococci concentrations suggests that a quicker turnaround of samples would assist with more responsive health risk communication. However, it may be argued that if there is very high rainfall across most days of the week, then the current regime of data collection and reporting is adequate. What our analysis and the current operational system cannot capture is high-flow events that are of very short duration (e.g. less than a day). However, people are unlikely to use these sites when flow is very high, i.e. exposure is low, suggesting that a time lag of 48 hours may not result in a large variation in the potential health risk. Evaluation of the timeliness of any environmental and public health surveillance systems depend on the purpose and goals of the surveillance system. Our findings provide a starting point to discuss the development of timely environmental early warning systems to protect public health.

Our limited prediction accuracy of high intestinal enterococci concentrations during low-flow conditions using water quality parameters combined with the high visitation rates to these locations during this time suggest a role of human sources of contamination including pets, children in nappies, and swimming stirring up sediments. [Elmir *et al.* \(2007\)](#) found that bathers transport a significant amount of intestinal enterococci and staphylococcus aureus into the water column. In Florida, management actions that included restricted pet access, bird management policies, and access fees were associated with a lower frequency of intestinal enterococci exceedances ([Kelly *et al.* 2018](#)). Intestinal enterococci does not identify the specific pathogens that are capable of causing human illness. The presence of high concentrations of FIB in water is indicative of faecal contamination, but due to their ubiquitous presence, concentrations alone do not provide information on the source(s) of contamination ([Codello *et al.* 2021](#)). For example, human faecal pollution typically presents the greatest risk because of the possible presence of human pathogens, while cattle manure may be a close second because of the possible presence of zoonotic pathogens such as *Cryptosporidium* spp. and enteropathogenic *Escherichia coli*. Microbial source tracking methods and environmental DNA (eDNA) barcoding can help identify sources of faecal contamination in complex environments ([Yan & Sadowsky 2007](#); [Staley *et al.* 2018](#)). When combined with different base flow conditions, such nuanced information can help inform management options to reduce non-point sources of contamination, while protecting public health.

Regression-based models such as the ones useful in operational contexts and for short-term predictions, but cannot be used to understand underlying processes, achieved mainly through the use of mechanistic models ([de Brauwere *et al.* 2014](#)). Another approach is to use a Bayesian framework similar to that used by [Farnham & Lall \(2015\)](#) to predict bacterial water contamination in New York City Harbour. This approach provides the predictions with uncertainty ranges included in model output, which can help water managers to better assess the risk of contamination to water users. Linking such macro-ecological data with hydrological and health outcome data could provide evidence-based solutions to contemporary public health problems, and in-turn help to develop locality-specific guidelines for water quality ([Verhoughstraete *et al.* 2020](#)) that will enhance aquatic ecosystem health.

Including rainfall and the full range of hydrological characteristics and other drivers of contamination and flow would be important in future investigations. For example, a recent study of flow exceedance in relation to urban land-use has shown that impervious surfaces correlated with increased frequency of high-flow events in Missouri streams ([Zeiger & Hubbard 2019](#)). Although including rainfall is a natural extension of the current work, a similar modelling study in Hawaii showed that in addition to rainfall, including the relative change in discharge incorporates a proximate representation of the energy available to transport particulates, improving predictions of near-shore water quality ([Strauch 2017](#)). It is important to note that rainfall itself may not be a useful predictor of intestinal enterococci concentrations, due to the complex hydrology of water catchments. [French \(2012\)](#) used multiple regression analyses of rainfall and intestinal enterococci concentrations in coastal waters of southern New South Wales, and found no statistical significance between the two variables, although we acknowledge the difference between coastal waters and inland rivers. So, while including rainfall is important, discharge and flow may be a better predictor of the proximate drivers of changing intestinal enterococci concentrations. Despite the limitations in the data and analytical framework used here, our results provide useful insights on high- and low-flow-related drivers of intestinal enterococci concentration across multiple sites over a long time period.

5. CONCLUSIONS

There is a need to monitor flow conditions for public health in natural, recreational swimming areas. Our findings, consistent across multiple sites over a long time period indicate that management actions that relate to high- and low-flow events leading to increased intestinal enterococci in these inland systems may have to be different, with implications for the health risks associated with natural swimming areas globally. With climate change projections indicating a greater likelihood of reduced rainfall (and thereby streamflow) in southeast Australia, risks associated with low streamflow and increased FIB could grow. Similarly, with the intensity of severe storm events likely to increase under climate change projections, the risks associated with the increased FIB concentrations under higher flows could also increase. As the risk is anticipated to increase, enhancing the ability to predict and model FIB concentrations would enhance the timeliness of risk communication and therefore decrease the public health risks to the community.

ACKNOWLEDGEMENTS

We thank staff from Icon water and ACT Parks and Conservation Service for helpful discussion that improved this manuscript. An anonymous reviewer provided constructive feedback that improved the manuscript. Funding for a research assistant to undertake this research was provided by the Australian National University's National Centre for Epidemiology and Population Health.

COMPETING INTERESTS

The authors have no competing interests to declare

DATA AVAILABILITY STATEMENT

Data cannot be made publicly available; readers should contact the corresponding author for details.

REFERENCES

- ABARES (Australian Bureau of Resource and Agricultural Economics and Sciences) 2021 *Catchment Scale Land Use Australia – Update December 2020 [WWW Document]*. Available from: <https://www.agriculture.gov.au/abares/aclump/catchment-scale-land-use-of-australia-update-december-2020> (accessed 5 January 2021).
- Abia, A. L. K., James, C., Ubomba-Jaswa, E. & Benteke Momba, M. N. 2017 *Microbial remobilisation on riverbed sediment disturbance in experimental flumes and a human-impacted river: implication for water resource management and public health in developing Sub-Saharan African countries*. *Int. J. Environ. Res. Public Health* **14**, 306. <https://doi.org/10.3390/ijerph14030306>.
- ACT Government Health 2014 *ACT Guidelines for Recreational Water Quality*. Canberra (accessed 17 December 2019).
- Alkolibi, F. M. 2002 *Possible effects of global warming on agriculture and water resources in Saudi Arabia: impacts and responses*. *Clim. Change* **54**, 225–245. <https://doi.org/10.1023/A:1015777403153>.
- Anderson, S. A., Turner, S. J. & Lewis, G. D. 1997 *Enterococci in the New Zealand environment: implications for water quality monitoring*. *Water Sci. Technol.* **35**, 325–331. [https://doi.org/10.1016/S0273-1223\(97\)00280-1](https://doi.org/10.1016/S0273-1223(97)00280-1).
- ANZECC & ARMCANZ 2000 *Australian and New Zealand Guidelines for Fresh and Marine Water Quality*. Australian and New Zealand Environment and Conservation Council and Agriculture and Resource Management Council of Australia and New Zealand, Canberra.
- Badgley, B. D., Nayak, B. S. & Harwood, V. J. 2010 *The importance of sediment and submerged aquatic vegetation as potential habitats for persistent strains of enterococci in a subtropical watershed*. *Water Res.* **44**, 5857–5866. <https://doi.org/10.1016/j.watres.2010.07.005>.
- Boehm, A. & Sassoubre, L. 2014 *Enterococci as indicators of environmental fecal contamination*. In: *Enterococci: From Commensals to Leading Causes of Drug Resistant Infection* (Glimore, M., Clewell, D., Ike, Y. & Al, E. eds.). Massachusetts Eye and Ear Infirmary, Boston.
- BOM (Bureau of Meteorology) & CSIRO (Commonwealth Scientific and Industrial Research Organisation) 2018 *State of the Climate 2018* (accessed 17 December 2019).
- Buckerfield, S. J., Quilliam, R. S., Waldron, S., Naylor, L. A., Li, S. & Oliver, D. M. 2019 *Rainfall-driven E. coli transfer to the stream-conduit network observed through increasing spatial scales in mixed land-use paddy farming karst terrain*. *Water Res. X* **5**, 100038. <https://doi.org/10.1016/j.wroa.2019.100038>.
- Cho, K., Pachepsky, Y., Kim, J. H., Guber, A., Shelton, D. & Rowland, R. 2010a *Release of Escherichia coli from the bottom sediment in a first-order creek: experiment and reach-specific modeling*. *J. Hydrol.* **391**, 322–332. <https://doi.org/10.1016/j.jhydrol.2010.07.033>.
- Cho, K. H., Cha, S. M., Kang, J.-H., Lee, S. W., Park, Y., Kim, J.-W. & Kim, J. H. 2010b *Meteorological effects on the levels of fecal indicator bacteria in an urban stream: a modeling approach*. *Water Res.* **44**, 2189–2202. <https://doi.org/10.1016/j.watres.2009.12.051>.

- Cho, K. H., Pachepsky, Y. A., Oliver, D. M., Muirhead, R. W., Park, Y., Quilliam, R. S. & Shelton, D. R. 2016 Modeling fate and transport of fecally-derived microorganisms at the watershed scale: state of the science and future opportunities. *Water Res.* **100**, 38–56. <https://doi.org/10.1016/j.watres.2016.04.064>.
- Chu, Y., Tournoud, M. G., Salles, C., Got, P., Perrin, J. L., Rodier, C., Caro, A. & Troussellier, M. 2014 Spatial and temporal dynamics of bacterial contamination in South France coastal rivers: focus on in-stream processes during low flows and floods. *Hydrol. Process.* **28**, 3300–3313. <https://doi.org/10.1002/hyp.9900>.
- Codello, A., McLellan, S. L., Steinberg, P., Potts, J., Scanes, P., Ferguson, A., Hose, G. C., Griffith, M., Roguet, A., Lydon, K. A. & Maher, W. A. 2021 A weight-of-evidence approach for identifying potential sources of untreated sewage inputs into a complex urbanized catchment. *Environ. Poll.* **275**, 116575.
- Coffey, R., Dorai-Raj, S., O'Flaherty, V., Cormican, M. & Cummins, E. 2013 Modeling of pathogen indicator organisms in a small-scale agricultural catchment using SWAT. *Hum. Ecol. Risk Assess. An Int. J.* **19**, 232–253. <https://doi.org/10.1080/10807039.2012.701983>.
- Curriero, F. C., Patz, J. A., Rose, J. B. & Lele, S. 2001 The association between extreme precipitation and waterborne disease outbreaks in the United States, 1948–1994. *Am. J. Public Health* **91**, 1194–1199. <https://doi.org/10.2105/AJPH.91.8.1194>.
- Dale, K., Kirk, M., Sinclair, M., Hall, R. & Leder, K. 2010 Reported waterborne outbreaks of gastrointestinal disease in Australia are predominantly associated with recreational exposure. *Aust. N. Z. J. Public Health* **34**, 527–530. <https://doi.org/10.1111/j.1753-6405.2010.00602.x>.
- Daly, E., Kolotelo, P., Schang, C., Osborne, C. A., Coleman, R., Deletic, A. & McCarthy, D. T. 2013 *Escherichia coli* concentrations and loads in an urbanised catchment: the Yarra River, Australia. *J. Hydrol.* **497**, 51–61. <https://doi.org/10.1016/j.jhydrol.2013.05.024>.
- Davis, J., O'Grady, A. P., Dale, A., Arthington, A. H., Gell, P. A., Driver, P. D., Bond, N., Casanova, M., Finlayson, M., Watts, R. J., Capon, S. J., Nagelkerken, I., Tingley, R., Fry, B., Page, T. J. & Specht, A. 2015 When trends intersect: the challenge of protecting freshwater ecosystems under multiple land use and hydrological intensification scenarios. *Sci. Total Environ.* **534**, 65–78. <https://doi.org/10.1016/j.scitotenv.2015.03.127>.
- de Brauwere, A., Ouattara, N. K. & Servais, P. 2014 Modeling fecal indicator bacteria concentrations in natural surface waters: a review. *Crit. Rev. Environ. Sci. Technol.* **44**, 2380–2453. <https://doi.org/10.1080/10643389.2013.829978>.
- Deely, J., Hodges, S., McIntosh, J. & Bassett, D. 1997 Enterococcal numbers measured in waters of marine, lake, and river swimming sites of the Bay of Plenty, New Zealand. *New Zeal. J. Mar. Freshw. Res.* **31**, 89–101. <https://doi.org/10.1080/00288330.1997.9516747>.
- Elmir, S. M., Wright, M. E., Abdelzaher, A., Solo-Gabriele, H. M., Fleming, L. E., Miller, G., Rybolowik, M., Peter Shih, M.-T., Pillai, S. P., Cooper, J. A. & Quaye, E. A. 2007 Quantitative evaluation of bacteria released by bathers in a marine water. *Water Res.* **41**, 3–10. <https://doi.org/10.1016/j.watres.2006.10.005>.
- French, C. E. 2012 *Concentrations of Enterococci in Relation to Rainfall in Sutherland Shire Catchments Concentrations of Enterococci in Relation to Rainfall in Sutherland Shire*. University of Wollongong (accessed 17 January 2021).
- Gao, G., Falconer, R. A. & Lin, B. 2013 Modelling importance of sediment effects on fate and transport of enterococci in the Severn Estuary, UK. *Mar. Pollut. Bull.* **67**, 45–54. <https://doi.org/10.1016/j.marpolbul.2012.12.002>.
- Herrig, I., Seis, W., Fischer, H., Regnery, J., Manz, W., Reifferscheid, G. & Böer, S. 2019 Prediction of fecal indicator organism concentrations in rivers: the shifting role of environmental factors under varying flow conditions. *Environ. Sci. Eur.* **31**, 59. <https://doi.org/10.1186/s12302-019-0250-9>.
- IPCC 2014 *Climate Change 2014: Synthesis Report*. Contribution of Working Groups I, II, and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Geneva (accessed 19 December 2019).
- Jamieson, R., Joy, D., Lee, H., Kostaschuk, R. & Gordon, R. 2005 Resuspension of sediment-associated in a natural stream. *J. Environ. Qual.* **34**, 581–589. <https://doi.org/10.2134/jeq2005.0581>.
- Joosten, R., Sonder, G., Parkkali, S., Brandwagt, D., Fanoy, E., Mughini-Gras, L., Lodder, W., Ruland, E., Siedenburg, E., Kliffen, S. & van Pelt, W. 2017 Risk factors for gastroenteritis associated with canal swimming in two cities in the Netherlands during the summer of 2015: a prospective study. *PLoS One* **12**, e0174732.
- Kelly, E. A., Feng, Z., Gidley, M. L., Sinigalliano, C. D., Kumar, N., Donahue, A. G., Reniers, A. J. H. M. & Solo-Gabriele, H. M. 2018 Effect of beach management policies on recreational water quality. *J. Environ. Manage.* **212**, 266–277. <https://doi.org/10.1016/j.jenvman.2018.02.012>.
- Kistemann, T., Claßen, T., Koch, C., Dangendorf, F., Fischeider, R., Gebel, J., Vacata, V. & Exner, M. 2002 Microbial load of drinking water reservoir tributaries during extreme rainfall and runoff. *Appl. Environ. Microbiol.* **68**, 2188–2197. <https://doi.org/10.1128/AEM.68.5.2188-2197.2002>.
- Lal, A. & Hargreaves, J. 2020 An epidemiologic approach to environmental monitoring: cyanobacteria in Australia's Murray–Darling basin. *Stoch. Environ. Res. Risk Assess.* **34** (7), 949–958.
- Meyer, J. L., Paul, M. J. & Taulbee, W. K. 2005 Stream ecosystem function in urbanizing landscapes. *J. North Am. Benthol. Soc.* **24**, 602–612. <https://doi.org/10.1899/04-021.1>.
- Olley, J. M. & Wasson, R. J. 2003 Changes in the flux of sediment in the upper Murrumbidgee catchment, Southeastern Australia, since European settlement. *Hydrol. Process.* **17**, 3307–3320. <https://doi.org/10.1002/hyp.1388>.
- O'Reilly, W., Brademann, A., Ferronato, B., Kellock, D., Lind, M. & Ubrihien, R. 2021 *Catchment Health Indicator Program: Report Card 2020*. Upper Murrumbidgee Waterwatch, Canberra.

- Ribolzi, O., Evrard, O., Huon, S., Rochelle-Newall, E., Henri-Des-Tureaux, T., Silvera, N., Thammahacksac, C. & Sengtaeuanghoung, O. 2016 Use of fallout radionuclides (7Be, 210pb) to estimate resuspension of *Escherichia coli* from streambed sediments during floods in a tropical montane catchment. *Environ. Sci. Pollut. Res.* **23**, 3427–3435. <https://doi.org/10.1007/s11356-015-5595-z>.
- Sidhu, J. P. S., Hodgers, L., Ahmed, W., Chong, M. N. & Toze, S. 2012 Prevalence of human pathogens and indicators in stormwater runoff in Brisbane, Australia. *Water Res.* **46**, 6652–6660. <https://doi.org/https://doi.org/10.1016/j.watres.2012.03.012>.
- Signor, R. S., Roser, D. J., Ashbolt, N. J. & Ball, J. E. 2005 Quantifying the impact of runoff events on microbiological contaminant concentrations entering surface drinking source waters. *J. Water Health* **3**, 453–468. <https://doi.org/10.2166/wh.2005.052>.
- Smith, A., Western, A. & Hannah, M. 2013 Linking water quality trends with land use intensification in dairy farming catchments. *J. Hydrol.* **476**, 1–12. <https://doi.org/10.1016/j.jhydrol.2012.08.057>.
- Staley, Z. R., Chuong, J. D., Hill, S. J., Grabuski, J., Shokralla, S., Hajibabaei, M. & Edge, T. A. 2018 Fecal source tracking and eDNA profiling in an urban creek following an extreme rain event. *Sci. Rep.* **8**, 14390. <https://doi.org/10.1038/s41598-018-32680-z>.
- Strauch, A. M. 2017 Relative change in stream discharge from a tropical watershed improves predictions of fecal bacteria in near-shore environments. *Hydrol. Sci. J.* **62**, 1381–1393. <https://doi.org/10.1080/02626667.2017.1310381>.
- Suter, E., Juhl, A. & O'Mullan, G. 2011 Particle association of enterococcus and total bacteria in the lower Hudson River estuary, USA. *J. Water Resour. Prot.* **3**, 715–725. <https://doi.org/0.4236/jwarp.2011.310082>.
- Terry, J., Benskin, C., Eastoe, E. & Haygarth, P. 2014 Temporal dynamics between cattle in-stream presence and suspended solids in a headwater catchment. *Environ. Sci. Process. Impacts* **16**, 1570–1577. <https://doi.org/10.1039/c3em00686g>.
- Thilakarathne, M., Sridhar, V. & Karthikeyan, R. 2018 Spatially explicit pollutant load-integrated in-stream *E. coli* concentration modeling in a mixed land-use catchment. *Water Res.* **144**, 87–103. <https://doi.org/https://doi.org/10.1016/j.watres.2018.07.021>.
- Tornevi, A., Axelsson, G. & Forsberg, B. 2013 Association between precipitation upstream of a drinking water utility and nurse advice calls relating to acute gastrointestinal illnesses. *PLoS One* **8**, e69918–e69918. <https://doi.org/10.1371/journal.pone.0069918>.
- Verhougstraete, M. P., Pogreba-Brown, K., Reynolds, K. A., Lamparelli, C. C., Zanolli Sato, M. I., Wade, T. J. & Eisenberg, J. N. S. 2020 A critical analysis of recreational water guidelines developed from temperate climate data and applied to the tropics. *Water Res.* **170**, 115294. <https://doi.org/10.1016/j.watres.2019.115294>.
- Walters, E., Schwarzwälder, K., Rutschmann, P., Müller, E. & Horn, H. 2014 Influence of resuspension on the fate of fecal indicator bacteria in large-scale flumes mimicking an oligotrophic river. *Water Res.* **48**, 466–477. <https://doi.org/https://doi.org/10.1016/j.watres.2013.10.002>.
- Webb, L. & Hennessy, K. 2015 *Climate Change in Australia: Projections for Selected Australian Cities*. CSIRO and Bureau of Meteorology, Canberra. ISBN: 978-1-4863-0531-5.
- Williamson, C. E., Madronich, S., Lal, A., Zepp, R. G., Lucas, R. M., Overholt, E. P., Rose, K. C., Schladow, S. G. & Lee-Taylor, J. 2017 Climate change-induced increases in precipitation are reducing the potential for solar ultraviolet radiation to inactivate pathogens in surface waters. *Sci. Rep.* **7**, 13033. <https://doi.org/10.1038/s41598-017-13392-2>.
- Yan, T. & Sadowsky, M. J. 2007 Determining sources of fecal bacteria in waterways. *Environ. Monit. Assess.* **129**, 97–106. <https://doi.org/10.1007/s10661-006-9426-z>.
- Zeiger, S. J. & Hubbart, J. A. 2019 Quantifying relationships between urban land use and flow frequency of small Missouri streams. *Sci. Total Environ.* **659**, 1008–1015. <https://doi.org/https://doi.org/10.1016/j.scitotenv.2018.12.416>.

First received 25 November 2021; accepted in revised form 16 March 2022. Available online 7 April 2022